
Initiated: 5/12/04

Reviewed/revised: 10/10/07

Revision: 1

MILWAUKEE COUNTY EMS MEDICAL PROTOCOL TRAUMA TEAM ACTIVATION PEDIATRIC PATIENTS

Approved by: Ronald Pirrallo, MD, MHSA Signature: Page 1 of 1



CRITERIA:

Patient with any of the following:

Airway:

Intubation/assisted ventilation

Breathing:

Respiratory arrest

Respiratory distress (ineffective respiratory effort, stridor or grunting

Respiratory Rate:

0 - 5 months: RR less than 20 6 months -12 years: RR less than 16 13 years or older: RR less than 12

Circulation:

0 - 5 months: Systolic BP less than 60 or Pulse less than 90

6 months - 5 years: Systolic BP less than 70 or Pulse less than 70

6 years or older: Systolic BP less than 80 or Pulse less than 60

Clinical signs of shock:

Pale, cold, clammy, tachycardia with weak pulses, capillary refill greater then 3 seconds in a warm environment

Patients receiving blood

Neurological:

GCS less than 13

Anatomic Diagnosis:

Penetrating injury to head, neck, torso, groin Flail chest

Open chest wound

<u>Level I Trauma</u>

CRITERIA:

Trauma patient with any of the following, who do not meet the Level I criteria:

Open or depressed skull fracture - blunt trauma

Suspected spine or spinal cord injury Bilateral femur fractures Pelvic fracture

Suspected airway injury from facial burns,

chemical or smoke inhalation Complete/partial amputaton or degloving about wrist or above ankle

Deep penetrating injuries above knee Significant blunt maxillofacial trauma

<u>↓</u>
<u>Level I I Trauma</u>

CRITERIA:

Patient with any of the following mechanisms of injury, who does not meet the Level I or Level II criteria:

Falls over 10 feet Crash speed more than 20 mph Passenger ejected from vehicle Vehicle roll-over

Death or severe injury of same-car occupant

Pedestrian hit by vehicle and thrown,

dragged or run over
Pedestrian struck at 20 mph or more
Passenger compartment intrusion

more than 12"
Prolonged extrication - over 20
minutes

ED AIII